

()

//

(Chrysobothris parvipunctata)

()

(.)

(*Punica granatum* L.)

()



()



)

(

()

Chrysobothris (Obenb.)

(*parvipunctata*)

()

(Lebert et Cohn)Schroet ()

Phytophthora cactorum

Phytophthora (Smith et Smith)Leonian ()

citrophthora



جهت تهیه فایل **WORD** این مقاله به سایت DaneshResan.com مراجعه نمایید و عنوان مقاله را جستجو کنید
بیش از ۲ میلیون مقاله فارسی در این سایت موجود میباشد



EC

MA PDA CMA

CMA

()

()

() :

()

pH · EC

*			
/ a			
/ b	/	/	+
/ b			()
	%		*

Pleuroplaconema

punicae Chaudhuri and Singh

()


F			
/ **	/	/	
	/	/	
	/	/	
%	**	/	:

Phytophthora citrophthora

Phytophthora cactorum

(*Meloidogyne* spp.)

Original Order	Ranked Order	
Mean 1=65.63	Mean 1=65.63a	
Mean 2=5.500	Mean 2=5.500b	+
Mean 3=1.500	Mean 3=1.500b	()



EC

pH

)
(

« »

()



()

()

(*Chrysobothris parvipunctata*)

REFERENCES

10. Coolen, J. & C.J. D'herde. 1972. A method for the quantitative extraction of eggs and second stage juveniles of *Meloidogyne* spp. from soil. Merelbeke Belgium, 36 pp.
11. Coolen, W. A. & C.J. D'Herde. 1972. A method for the quantitative extraction of nematodes from plant tissue, Merelbeke Belgium, 77 pp.
12. Jenkins, W. R. 1964. A rapid centrifugal-flotation technique for separating nematodes from soil. Plant Dis. Rep., 48(9):692.