

(M.D)

(M.D) *

(M.Sc)

-
-

:

.

.

.

:

(American society of anesthesiologists) ASA I II

:

- /

- /

SPSS

/ ± /

/

/ :

± /

/ I

/ .

II

/

/

,

	/	/		/
	/			
			(p< /)	
± / /			± /	
	SPO ₂			(p< /)
				/
	/ .			
				:
				:



[]

()

()

Rolly Kay .

[]

()

()

Hunt Taveau

[]

Bovet

[]



%

/

()

II I

()

I

SPSS



/ /

/ ± /

± /

ASA

/

II

/ I

/

/

/

- /

(

)

/

()

/

- /

)

/

(

/


/

(p < /)

()

/

/



p < /		/	/	
		/	/	
N.S			/	
			/	

p < /		/		
		/	/	
			/	
p < /			/	
			/	

/

/


/

(/)

(p < /)

± /

± /



/

/

.() (p< /)

(P< /)

/

(P< /)

/

.()

.(P< /)

/

/

/


/ ()

/



()

[]



.[]



/

/

/

[]

/

()

-

/

± /

± /

/

/

()

/



[1] Stone DJ, Gal TJ. Airway management. In: Miller RD, ed. Anesthesia. 5th ed. Philadelphia: Churchill-Livingstone, 2000. p. 1430-2.

() []

[3] Leuwer M, Zuzan O. Side effects of neuromuscular blocking drugs. European Society of Anaesthesiologists (ESA). 1999; 9Rc 1 (Refresher courses).

[4] Finucane BT, Santora AH. Principles of Airway Management, 2nd ed, St. Louis, Mosby, 1996. p.154-6.

[5] Stoelting RK, Miller RD. Basics of anesthesia. 3rd ed. London: Churchill Livingstone; 2000. p.91-6.

[6] Rushman GB, Davies NJ, Cashman JN, Editors. Lee's synopsis of Anesthesia. 12th Ed. New Delhi: Butterworth-Heinemann; 1999. p.188.

[7] Collins L, Prentice J, Vaghadia H. Tracheal intubation of outpatients with and without muscle relaxants. Can J Anaesth, 2000; 47(5):427-32.

[8] Jabbour-Khoury SI, Dabbous AS, Rizk LB, Abou Jalad NM, Bartelmaos TE, El-Khatib MF, et al. A combination of alfentanil-lidocaine-propofol provides better intubating conditions than fentanyl-lidocaine-propofol in the absence of muscle relaxants. Can J Anaesth, 2003; 50(2):116-20.

[9] Andel H, Klune G, Andel D, Felfernig M, Donner A, Schramm W, et al. Propofol without muscle relaxants for conventional or fiberoptic nasotracheal intubation: a dose-finding study. Anesth Analg, 2000; 91(2):458-61.

[10] Stevens JB, Vescovo MV, Harris KC, Walker SC, Hickey R. Tracheal intubation using alfentanil and no muscle relaxant: is the choice of hypnotic important? Anesth Analg, 1997; 84(6):1222-6.

[11] Hui JK, Critchley LA, Karmakar MK, Lam PK. Co-administration of alfentanil-propofol improves laryngeal mask airway insertion compared to fentanyl-propofol. Can J Anaesth, 2002; 49(5):508-12.

[12] Scheller MS, Zornow MH, Saidman LJ. Tracheal intubation without the use of muscle relaxants: a technique using propofol and varying doses of alfentanil. Anesth Analg, 1992; 75(5):788-93.

[13] Chohedri AH, Maghsoudi B, Mahmodi M. Propofol Alfentanil intubation; without muscle relaxant in young adult males. Professional Med J, 2004; 11(2):164-70.
