

()

*

(/ / : // :)

x x

(P<0.01)

(/) x
(/)

(P<0.05)

(P<0.05)

(P<0.01)

(P<0.05)

x

x

(Farid, 1988; Kashan, 1993;
Kashan et al, 2005; Miraei-Ashtiani et al., 2003;
Saatchi et al., 2005)

)
(
(Miraei-Ashtiani et al.,
(2003) Satchi et al. 2003)
(St Croix White)
(/)
(Godfrey & Weis, 2005)
(Farid, 1988)
(2002) Momani Shaker et al.
× ×
/ /
× (MM) ×
× (GG) ×
× (GM)
(MG) (2008) Vacca et al.
± / ×
/ ± /
(Unal et al., 2006; Kashan, 1993; Kashan et al.,
2005; Bunch et al., 2004; burke et al., 2003;
snowder et al., 2003)
(Pelibuey)
/ /
×
×
(Gutiérrez et al.,
2005)

($p < 0.0001$)

× ($p < 0.01$)

($P < 0.01$)

GM GG MG (Planimeter)

×

×

SAS
(GLM)

$$y_{ijk} = \mu + G_i + year_j +$$

$$a(age_{ijk} - \bar{age}_{...}) + b(w_{ijk} - \bar{w}_{...}) + e_{ijk}$$

(2002) Momani Shaker et al.

$$y_{ijk} = \mu + G_i + year_j + a(age_{ijk} - \bar{age}_{...}) + b(w_{ijk} - \bar{w}_{...}) + e_{ijk}$$

/ ± / ± /

(Godfrey & Weis, 2005) Phillips et al. 2005

()

MM	MG	GM	GG	
()	()	()	()	*
/ ± /	/ ± /	/ ± /	/ ± /	()
/ ± /	/ ± /	/ ± /	/ ± /	()
/ ± /	/ ± /	/ ± /	/ ± /	()
/ ^b ± /	/ ^a ± /	/ ^a ± /	/ ^a ± /	()
/ ^b ± /	/ ^a ± /	/ ^a ± /	/ ^a ± /	()
/ ^b ± /	/ ^{ab} ± /	/ ^a ± /	/ ^a ± /	()
/ ± /	/ ± /	/ ± /	/ ± /	()
/ ± /	/ ± /	/ ± /	/ ± /	(%)
/ ± /	/ ± /	/ ± /	/ ± /	(%)

.(p<0.05)

*

**

Godfrey & Weis (/ %) × (/ %) (2005) (2004) Gokdal et al. (Karakas) (Ile de France) Cloete et al. (Merino Landsheep) (SA Mutton Merino) (P<0.05) MG MM GM GG (p<0.01) × (±) El Fadili (/) (2001) et al. (GM / MM) (Timahdite) × (D'man) × (Merinos precoce) × (Gutie´rrez et al., 2005) (Momani) Shaker et al., 2002) (2005) Kashan et al. (/ %) ×

(mm)	(%)	()	
/ ^a _± /	/ ± /	/ ^a _± /	GM
/ ^b _± /	/ ± /	/ ^a _± /	GG
/ ^{ab} _± /	/ ± /	/ ^a _± /	MG
/ ^a _± /	/ ± /	/ ^b _± /	MM

(P<0.05)

GG MG

(2006) Cloete et al.

/ /)

/

Momani (2005) Godfrey & Weis (

(2002) Shaker et al.

(P<0.01)

(P<0.05)

/

.()

×

/ /

(2005) Kashan et al.

× ×

()

(p<0.05)

(p<0.01)

(2005) Gutierrez et al.

:
 .()
 ×
 / /)
 ()
 .(p<0.05) (/ /)
 Momani Shaker et al. (2002)
 / × (/) /
 / × (/)
 (/)
 (2004) Gokdal et al.
 .(/ ± / / ± /)

(P<0.05)

(p<0.05)

MG GM GG

MM

Godfrey & Weis

(2005)

() × ()

(P<0.01) × (p<0.01)

()	()	()	()	()	()
/ b _± /	/ b _± /	/ a _± /	/ a _± /	/ ± /	/ a _± /
/ a _± /	/ a _± /	/ b _± /	/ b _± /	/ ± /	/ ab _± /
/ a _± /	/ a _± /	/ b _± /	/ ab _± /	/ ± /	/ b _± /
/ b _± /	/ ab _± /	/ ab _± /	/ b _± /	/ ± /	/ b _± /

MM	MG	GM	GG	() **
/ ± /	/ ± /	/ ± /	/ ± /	(%)
/ ± /	/ ± /	/ ± /	/ ± /	*** ()
/ ± /	/ ± /	/ ± /	/ ± /	
/ ± /	/ ± /	/ ± /	/ ± /	
/ b _± /	/ a _± /	/ a _± /	/ a _± /	
/ ± /	/ ± /	/ ± /	/ ± /	

(p<0.05)

*

**

(%)	(%)	(%)	
/ ± /	/ ± /	/ ± /	GG
/ ± /	/ ± /	/ ± /	GM
/ ± /	/ ± /	/ ± /	MG
/ ± /	/ ± /	/ ± /	MM

×

(2005) Kashan et al. (Vacca et al., 2008)

REFERENCES

1. Bunch, T. D., Evans, R. C., Wang, S., Brennand, C. P., Whittier, D. R. & Taylor, B. J. (2004). Feed efficiency, growth rates, carcass evaluation, cholesterol level and sensory evaluation of lambs of various hair and wool sheep and their crosses. *Small Ruminant Research*, 52, 239–245.
2. Burke, J. M., Apple, J. K., Roberts, W. J., Boeger, C. B. & Kegley, E. B. (2003). Effect of breed-type on performance and carcass traits of intensively managed hair sheep. *Meat Science*, 63, 309-315.
3. Cloete, J. J. E., Cloete, S. W. P., Olivier, J. J. & Hoffman, L. C. (2007). Terminal crossbreeding of Dorper ewes to Ile de France, Merino Landsheep and SA Mutton Merino sires: Ewe production and lamb performance. *Small Ruminant Research*, 69, 28-35.
4. El Fadili, M., Michaux, C., Detilleux, J. & Leroy, P. L. (2001). Evaluation of fattening performances and carcass characteristics of purebred, first and second cross lambs between Moroccan Timahdite, D'man and improved meat rams. *Animal Science*, 72, 251-257.
5. Farid, A. (1989). Direct, maternal and heterosis effects for slaughter and carcass characteristics in three breeds of fat-tailed sheep. *Livestock Production Science*, 23, 137-162.
6. Godfrey, R. W. & Weis, A. J. (2005). Post-weaning growth and carcass traits of St. Croix White and Dorper × St. Croix White lambs fed a concentrate diet in the US Virgin Islands. *Sheep and Goat Research*, 20, 32-36.
7. Gokdal, O., Ulker, H., Karakus, F., Cengiz, F., Temur, C. & Handil, H. (2004). Growth, feedlot performance and carcass characteristics of Karakas and crossbred lambs (F1) (Ile de France x Akkaraman (G1) x Karakas) under rural farm conditions in Turkey. *South African Journal of Animal Science*, 34, 223-232.
8. Gutiérrez, J., Rubio, M. S., & Méndez, R. D. (2005). Effects of crossbreeding Mexican Pelibuey sheep with Rambouillet and Suffolk on carcass traits. *Meat Science*, 70, 1–5.
9. Kashan, N., (1993). Study of growth and carcass quality of Chaal and Zandi sheep breeds and crossbred lambs. *Iranian Journal of Agricultural Science*, 24, 47–63. (In Farsi).
10. Kashan, N. E. J., Manafi Azar, G. H., Afzalzadeh, A., & Salehi, A. (2005). Growth performance and carcass quality of fattening lambs from fat-tailed and tailed sheep breeds. *Small Ruminant Research*, 60, 267-271.
11. Macit, M., Karaoglu, M., Esenbuga, N., Kopuzlu, S. & Dayioglu, H. (2001). Growth performance of purebred Awassi, Morkaraman and Tushin lambs and their crosses under semi-intensive management in Turkey. *Small Ruminant Research*, 41, 177-180.

12. Miraei-Ashtiani, S. R., Noshary, A. R. & Moradi Shahrababak, M. (2003). Evaluation of the lamb and feed-lot performances of three crossbred and one purebred genotypes of Iranian fat-tailed sheep. In: *Proceedings of the British Society of Animal Science*, p. 146.
13. Momani Shaker, M., Abdullah, A. Y., Kridli, R. T., Blaha, J., Sada, I. & Sovjak, R. (2002). Fattening performance and carcass value of Awassi ram lambs, F1 crossbreds of Romanov × Awassi and Charollais × Awassi in Jordan. *Czech Journal of Animal Science*, 47, 429–438.
14. Phillips, W. A., Brown, M. A., Dolezal, H. G. & G. Q. Fitch. 2005. Feedlot performance and carcass characteristics of lambs sired by Texel, Romanov, St. Croix or Dorset rams from Polypay and St. Croix ewes. *Sheep and Goat Research*, 20, 11-16.
15. Saatchi, M., Miraei-Ashtiani, S. R. & Zare Shahneh, A. (2005). Comparison of growth and feedlot traits in Kordi crossbred and purebred lambs (Crossbreeding between some Iranian fat tailed breeds). In: *Proceedings of the British Society of Animal Science*, p. 133.
16. Snowden, G. D. & Duckett, S. K. (2003). Evaluation of the South African Dorper as a terminal sire breed for growth, carcass, and palatability characteristics. *Journal of Animal Science*, 81, 368–375.
17. Unal, N., Akcapinar, H., Atasoy, F. & Aytac, M. (2006). Some reproductive and growth traits of crossbred genotypes produced by crossing local sheep breeds of Kivircik x White Karaman and Chios x White Karaman in steppe conditions. *Archiv Tierzucht, Dummerstorf*, 49, 55-63.
18. Vacca, G., Carcangiu, M. V., Dettori, M. L., Pazzola, M., Mura, M. C., Luridiana, S. & Tillocam, G. (2008). Productive performance and meat quality of Mouflon × Sarda and Sarda × Sarda suckling lambs. *Meat Science*, 80, 326-334.