

()

*

(// : // :)

(%)
()

(Kafee et al., 2000)

(Kafee et al., 2000)

(Babalar & Pirmoradian, 2000)

(1995) Fukuyama et al. . (Bialczyk et al., 2007)
(Ca- zeolite) .

(2005) Gull et al. .

(1998) Siddigi et al.

(Siddigi et al., 1998)

()

(Barton et al., 2006)

(Dixon & Weed, 1998)

(1997) Kapetanios et al. .

(Delshad et al., 2006)

%

()

(1996) Challinor et al.

EC . ± / pH .

(mmol/L)	N	P	K	Ca	Mg	S	Cl
(N+)	/	/	/		/	/	
(N-)	/	/	/		/	/	/
(mg/L)	Na	Fe	Mn	Zn	Cu	Mo	B
(N)	(N+)	/	/	/	/	/	

(g/g) .
 (mm/day) . (/ mg NH₄/g z)
 (cm²/day) .
 (g/m²/day) .
 (mg/m²/day) . (Wang et al., 2006)

)
 ((WPCF,AWWA. Standard methods., 1989)

%)
 (Le Bot et al., 2001)

)
 (beril
 // //

kjdale Tecator Analyzer
 (1030)
 (Emami, 1996)

()
 (Leaf area meter model: DELTA-T,
 DEVICES LTD)

(Standard methods: 4500-NH₃-
 C,1989)

(Spectrometer. uv/vis, Perkinelmer Lambda 201
 USA.)

(Karimi & Azizi,
 :1997)

- 2. Specific Leaf Ratio(SLR) (m²/m²)
- 3. leaf Weight Ratio(LWR) (cm²/g)
- 4. Stem Elongation Ratio(SER)
- 5. Leaf Expansion Ratio(LER)
- 6. Crop Growth Ratio(CGR)
- 7. Net Assimilation Rate(NAR)

1. Leaf Area Index(LAI)

Tei et al.

(Wang et al., 2006)

(2002a&b)

SAS 6.12

(2000) Walker et al.

Excel

(Fukuyama et al., 1995)

LAI %
 LER () SLA ()
 ()

(Kutuk et al., 2004)

CGR () SER
 ()

(Gul et al., 2005; Maloupa et al.,

- .1999; Gent, 2004)

%

beril

(cm ² /g)	(cm ² /day)	(m ² /m ²)
/ a	1.6/.9a	2/26a*
11/4.0b	67/91b	2/22b

% *

beril

(g/g)	(cm ² /g)	(m ² /m ²)	(cm)
/ a	/ a	/ a	a / a*
/ a	/ a	/ b	/ a
/ a	/ a	/ C	/ b

+

(g/m ² /day)	(mg/m ² *day)	(cm ² /day)	(mm/day)	(cm ² /g)
/ a	/ a	/ a	/ a	/ a
/ a	/ a	/ b	/ a	/ b
/ b	/ a	/ c	/ b	/ b

% *

%

beril

beril	
(%)	
/ b*	(T1)
/ a	(T2)
	×
/ b	N ⁺ × T1
/ a	N ⁺ × T2
/ c	N ⁻ × T1
/ c	N ⁻ × T2
%	
T1	=N ⁻
T2	=N ⁺

EC

beril	
(ppm)	
/ a*	N ⁺ × ZNH ₄ +P
/ b	N ⁺ × Z+P
/ b	N ⁺ +P
/ c	N ⁻ × ZNH ₄ +P
/ d	N ⁻ × Z+P
/ d	N ⁻ +P
%	
=P	=N ⁻
=ZNH ₄ +P	=Z+P
	=N ⁺

beril

beril	
(ppm)	
/ b*	(T1)
/ a	(T2)
%	

()

beril


beril	
(ppm)	
/ a*	(N ⁺)
/ b	(N ⁻)
	+
/ a	
/ b	
/ b	
%	

()

...
 % /
 % /
 (1999) Harland et al.

Gull et al. (2005) Caballero et al.
 (2005) Mohammad et al. (2005)
 (1996) Challinor et al.

(1999) Maloupa et al. (2008) Ayan et al. .



beril

=P

=N-

=N+

=ZNH₄+P

=Z+P

beril

()	()
/ a +	/ a*
/ a	/ a
/ b	
%	*

LeBot et al. .

(2001)

()

REFERENCES

1. Ayan, S., Yahaoglu, Z., Gercek, V. & Sahin, A. (2008). Utilization of zeolite as a substrate for containerized oriental spruce (*Picea orientalis* L.(Link)) seedling propagation. *Acta Horticulture*, 779, 583-590.
2. Babalar, M., Pirmordian, M. (TR). (2000). Tree Fruit Nutrition (2th ed). University of Tehran press. (In Farsi).
3. Barton, L. & Comer, T.D. (2006). Irrigation and fertilizer strategies for minimizing nitrogen leaching from turfgrass. *Agricultural Water Management*. 80 (1-3), 160-175.
4. Bialczyk, J., Lechowski, Z., Dziga, D. & Meja, E. (2007). Fruit yield of tomato cultivated on media with bicarbonate and nitrate/ ammonium as the nitrogen source. *Journal of plant nutrition*, 30, 149-161.
5. Caballero, R. P., Gil, J., Benitez, C. & Gonzalez, J. L. (2005). The effect of adding zeolite to soils in order to improve the N-K nutrition of Olive trees. Preliminary results. *American Journal of Agricultural and Biological Sciences*, 2(1), 321-324.
6. Challinor, P. F., Fuller, M. P. & Parkinson, J. (1996). Growth and development of sweet pepper plants on unloaded and nutrient loaded clinoptilolite. In: *Proceedings of ISOSC, 9th International congress on soilless culture*, Naaldwijk, Netherlands. 105-122.
7. Delshad, M., Kashi, A. K. & Babalar, M. (2006). Possibility of Using Organic Media and Finding Suitable Nutrient Solution for Soilless Culture of Greenhouse Tomato. *Iranian Journal of Agricultural Science*, 37(2), (176-186). (In Farsi).
8. Dixon, J. B. & Weed, S. B. (1989). Minerals in soil environments. *Soil Science Society of America*, 1, 873-911.
9. Emami, E. (1996). *Methods of Plant Analysis*. Vol(1). pp:25-28. Soil and water research organization.
10. Fukuyama, T., Nonami, H., Katayama, K. & Hashimoto, Y. (1995). Improvement of hydroponic culture media by adding calcium-zeolite. *ISHS Acta Horticulturae*. 396, (115-122).
11. Greenwood, D. J., Gastal, F., Lemaire, G., Lemaire, a., Draycott, A., Millard, P. & Neeteson, J. (1991). Growth rate and %N of field grown crops: theory and experiments. *Annual Botany*, 67, 181-190.
12. Gent, M. P. N. (2004). Effect of nitrogen and potassium supply on yield and tissue composition of greenhouse tomato. *Acta Horticulturae*, 644, 369-376.
13. Gul, A., Erogul, D. & Ongun, A. r. (2005). Comparison of the use of zeolite and perlite as substrate for crisp-head lettuce. *Elsevier, Scientia Horticulturae*, 106, 464-471.
14. Harland, J., Lane, S. & Price, D. (1999). Further experiment with recycled zeolite as a substrate for the sweet pepper crop. *Acta Horticulturae*, 482, 187-194.
15. Hartman, P. L., Mill, H. A. & Jones, J. B. (1986). The influence of nitrate: Ammonium ratios on growth, fruit development and element concentration floradel tomato plant. *Scientia Horticulturae*, 111, 487-490.
16. Kafee, M., Zand, E., Kamkar, B., Shareefee, H. R. & Goldanee, M. (TR). (2000). *Plant physiology*. University of Mashhad press. (In Farsi).
17. Kapetanios, E. G & Ioizidou, M. (1992). Heavy metal removal by zeolite in tomato cultivation using compost. *Acta Horticulturae*, 302, 63-71.
18. Karimi, M. & Azizi, M. (TR). (1997). *Plant growth analysis*. University of Mashhad press. (In Farsi).
19. Kutuk, C., G. Cayci & Heng, L. K. (2004). Effect of increasing salinity and ¹⁵N-labelled urea levels on growth N uptake, and water use efficiency of young tomato plants. *Australian Journal of soil research*, 32, 345-351.
20. Le Bot, J., Jeannequin, B. & Fabre, R. (2001). Growth and nitrogen status of soilless tomato plants following nitrate withdrawal from the nutrient solution. *Annals of botany*, 88, 361-370.
21. Maloupa, M., Samartzidis, C., Couloumbis, P. & Komninou, A. (1999). Yield quality and photosynthetic activity of greenhouse grown 'Madelon' rose on perlite-zeolite substrate mixtures. *Acta Horticulturae*, 481, 97-99.
22. Mohammad, M. J., Karma, N. S. & Al-latai fed, N. K. (2005). Response of croton growing in a zeolite-containing substrate to different concentrations of fertilizer solution. *Communications in soil science and plant analysis*, 35, 2283-2297.
23. Pisanu, A., Manca, B., Mule, P., Chessa, F. & Meloni, S. (2007). Amended soils with natural zeolites: analysis of two-year tests on greenhouse tomato. *Acta Horticulture*, 747, 211-218.
24. Siddigi, M. Y., Kronzucker, H. J., Britto, D. T. & Glass, A. D. M. (1998). Growth of a tomato crop at reduced nutrient concentrations as a strategy to limit eutrophication. *Journal of plant nutrition*, 21, 1879-1895.
25. Tei, F., Benincasa, P. & Guiducci, M. (2002a). Critical nitrogen concentration in processing tomato. *Agronomy Journal*, 18, 45-55.
26. Tei, F., Benincasa, P. & Guiducci, M. (2002b). Effect of N availability on growth, N uptake, light interception and photosynthetic activity in processing tomato. In: *Acta Horticulturae*, 571, 209-216.

...

:

27. Walker, R. L., Burns, I. G. & Moorby, J. (2000). Responses of plant growth rate to nitrogen supply: a comparison of relative addition and N interruption treatments. *Oxford journals, Experimental Botany*, 52, (355), 309-317.
28. Wang, Y., Liu, S., Xu, Z., Heu, T., Chuan, S. & Zhu, T. (2006). Ammonia removal from leachate solution using natural Chinese clinoptilolite. *Journal of Hazardous Materials*, 136, 735-740.
29. WPCF, APHP, AWWA. (1989). *Standard methods for the examination of water and wastewater*. Methods 4500-NH₂-B: (pp. 4-115).
30. WPCF, APHP, AWWA. (1989). *Standard methods for the examination of water and wastewater*. Methods 4500-NH₃-C: (pp.4-117).

