

...

() Shabanpour .

(*Hypophthalmichthys molitrix*)

Tangestani .

Perez- Alonso *et al.*,)

()

(2003)

()

Tokur .

(Rezaei *et al.*, 2003)

()

-

(*Cyprinus carpio*)

-

(Park *et al.*, 1996)

Al-)

(Sheviklo, 2000)

(bulushi *et al.*, 2005)

-

()

Cakli (Schubring, 1999)

(*Sander*)

(*Sardina pilchardus*)

(°C)


lucioperca

() / /

(ANOVA)
LSD
SPSS

(P < /) (Zar, 1996)


Excel



(A, B) ± (a, b, c)

()

/ / / /




(A, B)

±
(a, b, c)

(PV)

()

(TBA)



(A, B)


±
(a, b, c)

()

()

(TBA)

(TBA)



)

(A, B)

±
(a, b, c, d)

(

(TVB-N)

TVB-N

/ / /


()

(Heme iron)

()

/ /
/


() /



(A, B)

±
(a, b, c)

()



(A, B)


±
(a, b, c)

(ppm)



()

...

()



()



/ / / /
 / / / /

(Toyomizu *et al.*, 1981)

(Ben- Gigirey *et al.*, 1999)

Tokur

()

(Oncorhynchus mykiss)

- Neamati ()
/ (Taşkaya *et al.*, 2003)

.(Hoke *et al.*, 2000) /
Dragoev *et al.*, 1998; Olafsdottir *et al.*,)
.(1997; Gomes *et al.*, 2003

Al-Bulushi ()

- () Neamati
)
(1995) Huss (

TBA
.(Aubourg *et al.*, 2002)

TVB-N .

Mazorra-)
.(Manzano *et al.*, 2000

(2000) Lakshmanan .

TVB-N .(Huss, 1994)

Huss,) ()

...

- TVB-N (1995; Ababouch *et al.*, 1996
-
Cakli *et al.*,)
() Taşkaya (2005
(Oncorhynchus mykiss)
- TVB-N .
Castro *et al.*,)
() Chytiri (2007
TVB-N
(Oncorhynchus mykiss)
/
Namulema () Aubourg
()

References

- Ababouch, L.H., Souibri, L., Rhaliby, K., Ouahdi, O., Battal, M., Busta, F.F., 1996. Quality changes in sardines (*Sardina pilchardus*) stored in ice and at ambient temperature. *Food Microbiology* 13(2), 123-132.
- Al-Bulushi, I., Kasapis, S., Al-Oufi, H., Al-Mamari, S., 2005. Evaluating the quality and storage stability of fish burgers during frozen storage. *Fisheries Science* 71(3), 648-654.

-
- AOAC. 2000. Official methods of analysis. 17th ed. Association of Official Analytical chemists, Procedure, Washington, DC.
- ASTM. 1969. Manual on Sensory Testing Methods. American Society for Testing and Materials, 1916 Race Street, Philadelphia, pa. 19103, pp. 34-42.
- Aubourg, S.P., Lehmann, I., Gallardo, J., 2002. Effect of previous chilled storage on rancidity development in frozen horse mackerel (*Trachurus trachurus*). Journal of the Science of Food and Agriculture 82(15), 1764-1771.
- Baker, J., 2005. Processing fish finger with bread crumbs. University Putra Malaysia Publications, 142 pp.
- Ben-Gigirey, B., Vieites Baptista De Sousa, J.M., Villa, T.G., Barros-Velazquez, J., 1999. Chemical changes and Visual Appearance of Albacore Tuna as Related to Frozen Storage. Journal of Food Science 64(1), 20-24.
- Bligh, E.G., Dyer, W.A., 1959. A rapid method of total lipid extraction and purification. Canadian Journal of Physiology and Pharmacology 37, 911-917.
- Cakli, S., Taskaya, L., Kisla, D., Celik, U., Altinel, C., Cadum, A., Kilinc, B., Haji, R., 2005. Production and quality of fish fingers from different fish species. European Food Research Technology 220, 526-530.
- Castro, F.A.F., Santana, H.M.P., Campos, F.M., Costa, N.M.B., Silva, M.T.C., Salaro, A.L., Franceschini, S.C.C., 2007. Fatty acid composition of three freshwater fishes under different storage and cooking processes. Food Chemistry 103(4), 1080-1090.
- Chytiri, S., Chouliara, I., Savvaidis, I.N., Kontominas, M.G., 2004. Microbiological, chemical and sensory assessment of iced whole and filleted aquacultured rainbow trout. Food Microbiology 21(2), 157-165.
- Clark, E.M., Mahoney, A.W., Carpenter, C.E., 1997. Heme and total iron in ready-to-eat chicken. Journal of Agriculture and Food Chemistry 45(1), 124-126.
- Dragoev, S.G., Kiosev, D.D., Danchev, S.A., Ioncheva, N.I., Genov, N.S., 1998. Study on the oxidative processes in frozen fish. Bulgarian Journal of Agricultural Science 4, 55-65.
- Egan, H., Krik, R.S., Sawyer, R., 1997. Pearsons Chemical Analysis of food 9: 609-634.
- Gomes, H.A., Silva, E.N., Nascimento, M.R.L., Fukuma, H.T., 2003. Evaluation of the 2-thiobarbituric acid method for the measurement of lipid oxidation in mechanically deboned gamma irradiated chicken meat. Food Chemistry 80, 433-437.
- Hoke, M.E., Jahncke, M.L., Silva, J.L., Hearsbereger, J.O., Chamul, R.S., Suriyaphn, O., 2000. Stability of washed frozen mince from channel catfish frames. Journal of food science 65(6), 1083-1086.
- Huss, H.H., 1994. Assurance of seafood quality. Rome: FAO Fisheries Technical Paper, No. 334. 169 pp.
- Huss, H.H., 1995. Quality and quality changes in fresh fish. Rome: FAO Fisheries Technical Paper No. 348. 195 pp.
- Lakshmanan, P.T., 2000. Fish spoilage and quality assessment. In T. S. G. Iyer, M. K. Kandoran, Mary Thomas, & P. T. Mathew (Eds.), Quality assurance in seafood processing. (pp. 26-40). Cochin: Society Fisher Techno (India).
- Luo, Y.K., Kuwahara, R., Kaneniwa, M., Murata, Y., Yokoyama, M., 2001. Comparison of gel properties of surimi from alaska pollock and three freshwater fish species : Effects of thermal processing and protein concentration. Journal of Food Science 66(3), 548-554.
- Mazorra-Manzano, M.A., Pacheco-Aguilar, R., Díaz-Rojas, E.I., Lugo-Sánchez, M. E., 2000. Postmortem changes in black Skipjack muscle during storage in ice. Journal of Food Science 65, 774-779.
- Namulema, A., Muyonga, J.H., Kaaya, A.N., 1999. Quality deterioration in frozen Nile perch (*Lates niloticus*) stored at -13 and -27 °C. Food Research International 32(2), 151-156.
- Neamati, M., Shabanpour, B., Shabani, A., Gholizadeh, M., 2009. The effect of cold storage on lipid quality and sensory evaluation of fish burgers made from Carp (*Cyprinus carpio*) surimi and red meat. Journal of Agriculture Science and Natural Resources 16, 108-118.
- Olafsdottir, G., Martinsdottir, E., Oehlenschläger, J., Dalgaard, P., Jensen, B., Undeland, I., 1997. Methods to evaluate fish freshness in research and industry. Trend Food Science and Technology 8, 258-265.
- Park, S., Brewer, M.S., Novakofski, J., Bechtel, P.J., McKeith, P.K., 1996. Process and characteristics for a Surimi – like material made from beef and pork. Journal of Food Science 61, 422-427.
- Pérez-Alonso, F., Arias, C., Aubourg, S.P., 2003. Lipid deterioration during chilled storage of Atlantic pomfret (*Brama brama*). European Journal of Lipid Science and Technology 105(11), 661-667.

...

-
- Rezaei, M., Sahari, M.A., Moeini, S., Safari, M., Ghaffari, F., 2003. Comparison quality oil of anchovy Kilka in two methods of transport and cold storage. *Iranian Journal of Fisheries Sciences* 3, 97-108.
 - Schubring, R., 1999. DSC Studies on deep frozen fishery products. *Journal of Thermochemica Acta* 337: 89-95.
 - Shabanpour, B., Asgharzadeh, A., Hosseini, H., Abbasi, M., 2008. Lipid Quality Changes of Silver Carp (*Hypophthalmichthys molitrix*) during Frozen Storage. *Journal of Agriculture Science and Natural Resources* 15(1), 17-24.
 - Sheviklo, G.H., 2000. Paste production guide and fish paste products. Naghshe mehr publications, 2-12 pp.
 - Tangestani, R., Alizadeh doughikolaee, E., Elyasi, A., 2010. An analysis of the Organoleptic properties of three types of fish fingers produced from silver carp flesh. *Journal of Fisheries (Iranian Journal of Natural Resources)* 63 (1), 1-10
 - Taşkaya, L., Çaklı, Ş., Kışla, D., Kılınç, B., 2003. Quality Changes of Fish Burger from Rainbow Trout during Refrigerated Storage. *E.U. Journal of Fisheries and Aquatic Sciences* 20, 147-154.
 - Tokur, B., Ozkutuk, S., Atici, E., Ozyurt, G., Ozyurt, C.E., 2006. Chemical and sensory quality changes of fish fingers, made from mirror carp (*Cyprinus carpio* L.,1758), during frozen storage (-18 C). *Food Chemistry* 99, 335-341.
 - Toyomizu, M., Hanaoka, K., Yamaguchi, K., 1981. Effect of release of free acids by enzymatic hydrolysis of phospholipids on lipid oxidation during storage of fish muscle at -5 °C. *Bull Jap Soc Sci Fish* 47, 610-615.
 - Zar, J.H., 1996. *Biostatistical Analysis*. Third edition. Prentice Hall. Upper Saddle River, NJ, USA, 662pp.

Quality of Fish Finger Produced from Common carp (*Cyprinus carpio*) During Storage time at 4 °C

Sh. Hasani¹, E. Alizadeh doughikollae^{*1}, E. Hayati jafar beygi¹ and M. Kamyab yeghaneh¹

¹Department of Fisheries, Faculty of Natural Resource, Zabol University, I.R. Iran

(Received: 22/11/2011, Accepted: 17/03/2012)

Abstract

Fish finger is a semi-prepared product of fish that many studies have been done to improve its texture, taste, flavor and color. The present study aims at investigating the quality and sensory characteristics of fish fingers produced from fillet and surimi of common carp during storage in refrigerator. Fish fingers were packed and then stored in refrigerator (4 °C). Chemical parameters (fat content, heme iron, peroxide value (PV), Thiobarbituric Acid (TBA), total volatile basic nitrogen (TVB-N) were measured and sensory analysis (texture, smell, taste, color and total acceptance) were determined by panelists on 0, 3, 6 and 9 days. The amount of peroxide (meq O₂/kg fat fish), thiobarbituric acid (mg malondialdehyde/kg fish flesh) and total volatile bases nitrogen (mg N/100 g fish flesh) of fish fingers between treatments were significantly different (P< 0.05). While the values of fillet and surimi fish finger reached to 4.07, 0.9, 33.17 and 3.53, 0.77 and 13.73 at the end of storage, respectively. The increase of values in fillet fish fingers was more than surimi fish fingers (P< 0.05). Heme iron content in both treatments showed a significant decrease. Total fat of fillet fish finger showed a significant decrease in comparison to surimi at the end of storage. Results of sensory analysis indicated significant reduction in sensory characteristics of fish fingers. Based on sensory analysis of two different types of common carp fish fingers shelf life was determined six days in refrigerator.

Keywords: Chemical parameters, Common carp (*Cyprinus carpio*), Fish finger, Sensory analysis, Shelf life.