

(Oncorhynchus mykiss)

*

(/ / : / / :)

() MIC .

:

...

(Miranda

.and Zemelman, 2002)

(Uhland and

MIC

.Higgins, 2006)

/

MIC

.(Cipriano, 2000)

.(Janda and Abbott, 1996)

.(Miranda and Zemelman, 2002)

Akinbowale .

.(Akinbowale *et al.*, 2007)

(Smith *et al.*,

(Vivekanandhan *et al.*, 2002)

MIC .1994)

.(Miranda and Zemelman, 2002)

¹TSA

cfu/ml

TSA

(NCCLS, 1997)

Urea, Catalas,
Oxides, MR-VP, Nitrate, MIS, Glocuze, Simmon
citrate, TSI, OF

)

(

: (MIC)

MIC

[National Committee for Clinical
Laboratory Standards(NCCLS), 1997]

(Whitman and Mac Nair,

2004)

)

(

/)

/

(/

(cfu/ml)

1

urea	Catalase	Oxidase	VP	MR	Nitrate	Motility	Indol	H S	Glocuze	Simmon citrate	TSI	of	Of ⁺
	+	+	+		+	+		+	+		/	+	+
				+							/	+	+
	+		+	+	+		+		+	+	/	+	+
1	+	+	+		+	+	+	+	+	+	/	+	+

urea	Catalase	Oxidase	VP	MR	Nitrate	Motility	Indol	H S	Glocuze	Simmon citrate	TSI	of	Of ⁺
	+	+		+	+				+	+	/	+	+

urea	Catalase	Oxidase	VP	MR	Nitrate	Motility	Indol	H S	Glocuze	Simmon citrate	TSI	of	Of ⁺
	+	+	+	+	+	+			+	+	/	+	+
	+	+	+	+	+				+	+	/	+	+
	+	+	+	+	+	+	+		+	+	/	+	+
	+	+	+		+				+	+	/	+	+

urea	Catalase	Oxidase	VP	MR	Nitrate	Motility	Indol	H S	Glocuze	Simmon citrate	TSI	of	Of ⁺
+	+	+	+	+	+	+		+	+	+	/	+	+
+	+			+	+			+	+		/	+	+
+	+			+	+			+	+	+	/	+	+
+	+	+	+	+	+	+		+	+		/	+	+

urea	Catalase	Oxidase	VP	MR	Nitrate	Motility	Indol	H S	Glocuze	Simmon citrate	TSI	of	Of ⁺
+	+	+	+	+	+	+		+	+		/		+
+	+	+	+	+	+	+	+	+	+	+	/		+
+	+			+	+					+	/		+
+	+			+	+					+	/		+
+	+	+	+	+	+	+	+	+	+		/		+


urea	Catalase	Oxidase	VP	MR	Nitrate	Motility	Indol	H S	Glocuze	Simmon citrate	TSI	of	Of ⁺
+	+	+	+	+	+	+	+	+	+	+	/	+	+
+	+			+	+	+	+	+			/	+	+
											/	+	+
+	+	+	+	+	+	+			+	+	/	+	+

..

urea	Catalase	Oxidase	VP	MR	Nitrate	Motility	Indol	H S	Glucose	Simmon citrate	TSI	of	Of
	+	+	+	+	+	+			+		/	+	+

)

.(



(2005) Hatha

(Cipriano, 2000)

Miranda and Zemelman (2002)

(Uhland and Higgins,
/ MIC .2006)

(Miranda and Zemelman,2002)
bacilli
MIC Spangaard ()
MIC

efflux MIC

(Miranda and
Zemelman,2002)

,MIC
() Akinbowal

(Miranda and
(1995) Kerry Zemelman,2002)

/

[

...

References

- Akinbowale, O.L., Peng, H., Grant, P., Barton, M.D., 2007. Antibiotic resistance in bacteria isolated from three freshwater fish farms and an unpolluted stream in Denmark. *Antimicrobial Agents*. 30: 177-182
- Cipriano, R. C., 2001. *Aeromonas Hydrophila* and Motile *Aeromonads* Cepticemias of Fish. FISH Disease Leaflet.
- Hatha, M., Vivekanandhan, A.A., Cheristol, G.J., 2005. Antibiotic resistance pattern of motile *Aeromonads* from farm raised fresh water fish. 98:131-134.
- Janda, J.M., Abbott, S. L., 1996. Human pathogens, p. 151-173. *In* Austin, B., Altwegg, M., Gosling, P. J., Joseph, S. (ed.), *The genus Aeromonas*. John Wiley & Sons, New York, N.Y.
- Kerry, J., Hiney, M., Coyne, R., NicGabhainn, S., Gilroy, D., Cazabon, D., Smith, P., 1995. Fish feed as a source of oxytetracycline-resistant bacteria in the sediments under fish farms. *Aquaculture*. 131:101-113 .
- Miranda, C.D., Zemelman, R., 2002. Bacterial resistance to oxytetracycline in Chilean salmon farming. *Aquaculture*, 212:31-47.
- National Committee for Clinical Laboratory Standards (NCCLS) (2002) Performance standards for antimicrobial disk and dilution susceptibility tests for bacteria isolated from animals; Approved standard – 2nd ed. NCCLS document M31-A2. NCCLS, 940 West Valley Road, Suite 1400, Wayne, PA 19087, USA.
- Smith, P., Hiney, M.P., Samuelsen, O.B., 1994. Bacterial resistance to antimicrobial agents used in fish farming : a critical evaluation of method and meaning. *Pergamon*, 4:273-313.
- Uhland, F.C., Higgins. R., 2006. Evaluation of the susceptibility of *Aeromonas salmonicida* to oxytetracycline and tetracycline using antimicrobial disk diffusion and dilution susceptibility tests. *Aquaculture*. 257: 111-117.
- Vivekanandhan, G., Savithamani, K., Hatha, A.A.M., , P., 2002. Antibiotic resistance of *Aeromonas hydrophila* isolates from marketed fish and prawn of south India. *International Journal of Food Microbiology*. 76:165-168.
- Withman, K.A., Mac NAair, N.G., 2004. *Finfish and Shellfish Bacteriology Manual*. Iowa state press.

Evaluation of Oxytetracycline Resistance in *Aeromonas hydrophila* Isolated from Reared *Oncorhynchus mykiss*

S. Zarghami¹, A. Mirvaghefi*¹, H. Farahmand¹, M.A. Nematollahi¹, M. Mosavi¹ and B. Khalili¹

¹ Department of Fisheries, Faculty of Natural Resources, University of Tehran, I.R. Iran
(Received: 17/04/2011, Accepted: 22/01/2012)

Abstract

The intensive use of antimicrobial agents worldwide for prophylactic and therapeutic purposes has been associated with increase of bacterial resistance in the exposed microbial environment. Oxytetracycline is the most frequently used antibacterial in the cold water fish industry because of its broad spectrum of activity and low cost. *Aeromonas hydrophila* and other motile aeromonads are among the most common bacteria in freshwater habitats throughout the world, and these bacteria frequently cause disease among cultured fishes. In this study, 50% of isolates from intestine of rainbow trout were identified as *Aeromonas hydrophila* by using biochemical technique. Oxytetracycline MIC_s were determined using agar dilution method at dosage 0,1,2,4,8,10 µl/ml. % of *Aeromonas hydrophila* was resistant to 30 µg/ml of oxytetracycline. The use of oxytetracycline in aquaculture might produce some negative impact on the treatment of human infections as a consequence of either direct transmission of resistant pathogens to humans, or indirectly through the transfer of resistance genes from environmental bacteria to human pathogens which is major risk for public health.

Keywords: Antibiotic resistance, *Oncorhynchus mykiss*, *Aeromonas hydrophila*