

(Oncorhynchus mykiss)

*

(/ / : / / :)

± / *(Oncorhynchus mykiss)*

...

Morais *et al.*,)

(2007).

Tidwell and Allen)

(2002).

Cahu *et al.*,)

(Cahu *et al.*, 2003).

(2009).

Kanazawa *et al.*, 1985a,b ; Tocher *et*)

Coutteau *et al.*,) (al., 2003

Buchet *et al.*,) (1997

2000 ; Cahu *et al.*, 2003; Gisbert *et al.*, 2005 ;

(Morais *et al.*, 2007; Hamza *et al.*, 2008

et al., 2000)

(Koven *et al.*, 1993)

(Fontage

(Cahu *et al.*, 2003).

Izquierdo *et al.*, 2000 ;)

(Wold *et al.*, 2007

()

(Geurden *et al.*, 2006).

...

(Kanazawa *et al.*, 1981).

(Tocher *et al.*, 2008)

(Dabrowski *et al.*, 2007)

(mg/l) ± /
pH= / / (mg/l) /
/ (mg/l)

)
(:
)
(Merck BDH
()
± /

)
(
()
/ / /
/ /

...

()

(Kjeltec

Analyzer Unit 2300)

/
Soxtec) FOSS

(2050

(SGR) (FCR)
(CF) (WG%)

= (FCR)¹

() / ()

(AOSC, 1997)

× () / = (CF)²

() ()

Ln Ln) = (SGR)³

(

× / (

] = (WG)⁴

×() /[() ()

) = (Survival)

×(/

(Metcalf *et al.*, 1966)

BF₃ %

(Izquierdo *et al.*, 2002)

Varian (GC)

SGM 120m×0/25mm)

i.d., film thickness 0/25mm) BPX70

FID

¹ Food conversion ratio (FCR)

² Condition factor (CF)


³ Specific growth rate (SGR)

⁴ weight gain (WG)

(Supel Co. Fame Mix, USA)

$(p > /)$ $p /)$ $(p > /)$ $(<$
 $(p < /)$ $(p < /)$
 $(p > /)$ $p /)$ $(>$
 $(p < /)$
 $(p > /)$ $(p > /)$
 $(p > /)$ $(p < /)$
 $(p > /)$

$/ \pm /^{ab}$	$/ \pm /^{bc}$	$/ \pm /^c$	$/ \pm /^{ab}$	$/ \pm /^{ab}$	$/ \pm /^a$	$/ \pm /^a$
$/ \pm /^{ab}$	$/ \pm /^{bc}$	$/ \pm /^c$	$/ \pm /^{ab}$	$/ \pm /^{ab}$	$/ \pm /^a$	$/ \pm /^a$
(SGR)						
$/ \pm /^{ab}$	$/ \pm /^{ab}$	$/ \pm /^a$	$/ \pm /^c$	$/ \pm /^{abc}$	$/ \pm /^{abc}$	$/ \pm /^{bc}$
(FCR)						
$/ \pm /^{ab}$	$/ \pm /^{bc}$	$/ \pm /^c$	$/ \pm /^{ab}$	$/ \pm /^{ab}$	$/ \pm /^a$	$/ \pm /^a$
$/ \pm /$	$/ \pm /$	$/ \pm /$	$/ \pm /$	$/ \pm /$	$/ \pm /$	$/ \pm /^{n.s}$
$/ \pm /^b$	$/ \pm /^a$	$/ \pm /^a$	$/ \pm /^a$	$/ \pm /^a$	$/ \pm /^a$	$/ \pm /^a$
$(p < /)$				$(\pm \quad n=)$		



(= n=)

Kanazawa *et al.*,)

(1985b

-

(Gisbert *et al.*, 2005)

(Geurden *et al.*, 1995,1997)

(Kanazawa *et al.*,1981)

Izquierdo *et al.*, 2000;)

-

(Wold *et al.*, 2007

(*Cyprinus carpio*)

(*Scophthalmus maximus*)

(*Plecoglossus altivelus*)

(Dabrowski *et al.*, 2007; Abedian *et al.*, 2011)

n

n

n

(Olsen *et al.*, 1999)

n

n

-

de vono

(Tocher *et al.*, 2008)

Caballero *et al.*,)

HUFA .(2003

HUFA

HUFA

Sargent *et al.*,) HUFA n-

(Tago *et al.*, 1999) HUFA n- .(1999

DHA

DHA

.(Gisbert *et al.*, 2005)

: n) (HUFA) HUFA n-

(EPA, C : n) (ARA, C

.(Cahu *et al.*, 2003) A

Tocher *et al.*,)

.(2008 *Penaeus* (*Plecoglossus altivelis*)

Kanazawa *et al.*, 1981, 1985) (*japonicus*

.(a,b

HUFA

.(Ishikawa, 2007)

HUFA

Kanazawa *et al.*, 1981;

(Geurden *et al.*, 1995)

(*Plecoglossus altivelis*)

(*Cyprinus carpio*) Knife jaw

References

- Abedian, A. M., Sotoudeh, E., Rezaei, M.H., 2011. Dietary soybean phosphatidylcholine affects growth performance and lipolytic enzyme activity in Caspian brown trout (*Salmo trutta Caspius*) alevin. *Aquaculture Research* (42): 655-663.
- AOAC (Association of Official Analytical Chemists). 1995: Association of official analytical chemists, official methods of analysis, 16th edition. Arlington, VA, USA.
- AOCS (2009) Official Methods and Recommended Practices of the AOCS. 6th edition. American Oil Chemists' Society, Champaign, IL, USA.
- Buchet, V., Zambonino-Infante J. L., Cahu, C.L., 2000. Effect of lipid level in a compound diet on the development of red drum (*Sciaenops ocellatus*) larvae. *Aquaculture* (184), 339-347.
- Caballero, M. J., Izquierdo, M.S., Kjorsvik, E., Montero, D., Socorro, J., Fernandez, A.J., Roselund, G., 2003. Morphological aspects of Intestinal cells from gilthead Seabream (*Sparus aurata*) fed diets containing different lipid sources, *Aquaculture* (225), 325–340.
- Cahu, C. L., Zambonino-Infante, J. L., Barbosa, V., 2003. Effect of dietary phospholipid level and phospholipid: neutral lipid value on the development of sea bass (*Dicentrarchus labrax*) larvae fed a compound diet. *British Journal of Nutrition* (90), 21-28.
- Cahu, C. L., Gisbert, E., Villeneuve, L. A. N., Morais, S., Hamza, N., Wold, P.A., Zambonino-Infante, J.L., 2009. Influence of dietary phospholipids on early ontogenesis of fish. *Aquaculture Research* (40), 989-999.
- Coutteau, P., Geurden, I., Camara, M. R., Bergot, P., Sorgeloos, P., 1997. Review on the dietary effects of phospholipids in fish and crustacean larviculture. *Aquaculture* (155), 149–164.
- Dabrowski, K., Rinchar, J., Czesny, S., 2007. Influence of lipid class and fatty acid deficiency on survival, growth, and fatty acid composition in rainbow trout juveniles. *Aquaculture* (264), 363–371.
- Fontagne, S., Burtaire, L., Corraze, G., Bergot, P., 2000. Effects of mediumchain triacylglycerols (tricaprylin and tricaproin) and phospholipid supply on survival, growth and lipid metabolism in common carp (*Cyprinus carpio*) larvae. *Aquaculture* (190), 289–303.
- Geurden, I., Radunz-Neto, J., Bergot, P., 1995. Essentiality of dietary phospholipids for carp (*Cyprinus carpio L.*) larvae. *Aquaculture* (131), 303–314.
- Geurden, I., Coutteau, P., Sorgeloos, P., 1997. Effect of dietary phospholipid supplementation on growth and fatty acid composition of European sea bass (*Dicentrarchus labrax L.*) and turbot (*Scophthalmus maximus L.*) juveniles from weaning onwards. *Fish Physiology and Biochemistry* (16), 259–272.
- Geurden, I., Gondouin, E., Rimbach, M., Koppe, W., Kaushik, S., Boujard, T., 2006. The evaluation of energy intake adjustments and preferences in juvenile rainbow trout fed increasing amounts of lipid. *Physiology and Behavior* (88), 325-332.
- Gisbert, E., Villeneuve, L., Zambonino-Infante, J.L., Quazuguel, P., Cahu, C.L., 2005. Dietary Phospholipids are more efficient than neutral lipids for long-chain polyunsaturated fatty acid supply in european sea Bass (*Dicentrarchus labrax*) Larval Development. *Lipids* (40), 609-618.
- Hamza, N., Mhetli, M., Khemis, I.B., Cahu, C., Kestemont, P., 2008. Effect of dietary phospholipids levels on performance, enzyme activities and fatty acid composition of pikeperch (*Sander lucioperca*) larvae. *Aquaculture* (275), 274–282.
- Institute of Standards and Industrial Research of Iran, Animal and vegetable fats and oils, Determination of phosphorus content-Part 1: Colorimetric method, number 10741-1. 1-st. edition (in persian).
- Ishikawa, M., 2007. Lipids. In: Dietary supplements for the health and quality of cultured fish (Nakagawa, H., Minoru, S and Delbert, M. G., eds), 64-73. Crown press. UK.
- Izquierdo, M. S., Socorro, J., Arantzamendi, L., Hernandez-Cruz, C.M., 2000. Recent advances in lipid nutrition in fish larvae. *Fish Physiology and Biochemistry* (22), 97-107.
- Izquierdo, M., Liu, J., Caballero, M.J., Ali, T.E.S., Hernandez-cruz, C. M., Valencia, A., Fernandez-palacios, H., 2002. Necessity of dietary lecithin and eicosapentaenoic acid for growth, survival, stress resistance and lipoprotein formation in gilthead seabream (*Sparus aurata*). *Fisheries Science* (68), 1165-1172.
- Kanazawa, A., Teshima, S., Inamori, S., Iwashita T., Nagao, A., 1981. Effects of phospholipids on survival rate and incidence of malformation in the larval ayu. Member Faculty of Fisheries Kagoshima University (30),301–309.

-
- Kanazawa, A., Teshima, S., sakamoto, M., 1985a. Effect of dietary lipids, fatty acids and phospholipids on the growth and survival of prawn (*Penaeus japonicas*) larvae. *Aquaculture* (50), 39-40
 - Kanazawa, A., Teshima, S., Sakamoto, M., 1985b. Effects of dietary bonito-egg phospholipids and some phospholipids on growth and survival of the larval ayu (*Plecoglossus altivelis*). *Journal of Applied Ichthyology* (4),165-170.
 - Koven, W.M., Kolkovski, S., Tandler, A., Kissil, G.W., Sklan, D., 1993. The effect of dietary lecithin and lipase, as function of age, on n-9 fatty acid incorporation in the tissue lipids of *Sparus aurata* larvae. *Fish Physiology and Biochemistry* (10), 357-364.
 - Metcalfe, L.D., Schmitz, A.A., Pelka, J. R., 1966. Rapid preparation of fatty acid esters from lipids for gas chromatographic analysis. *Analytical Chemistry* (38), 524-535.
 - Morais, S., Conceicao, I.E.C., Rønnestad, I., Koven, W., Cahu, C., Zambonino-Infante, J.L., Dinis, M.T., 2007. Dietary neutral lipid level and source in marine fish larvae: Effects on digestive physiology and food intake. *Aquaculture* (268), 106-122.
 - National Research Council (NRC), 1993. *Nutrient Requirements of Fish*. National Academy Press, Washington., 114 pp.
 - Olsen, R. E., Myklebust, R., Kaino, T., Ringo, E., 1999. Lipid digestibility and ultrastructural changes in the enterocytes of arctic char (*Salvelinus alpinus* L.) Fed Linseed Oil and Soybean Lecithin, *Fish Physiology and Biochemistry* (21), 35-44.
 - Sargent, J., Bell, G., McEvoy, L., Tocher, D., Estevez, A., 1999. Recent developments in the essential fatty acid nutrition of fish. *Aquaculture* (177),191-199.
 - Tago, A., Yamamoto, Y., Teshima, S., Kanazawa, A., 1999. Effects of 1,2-di-20:5 phosphatidylcholine (PC) and 1,2-di-22:6-PC on growth and stress tolerance of Japanese flounder (*Paralichthys olivaceus*) larvae. *Aquaculture* (179), 1-4.
 - Tidwell, J.H. Allan, G.L., 2002. Fish as food: aquaculture's contribution. *World Aquaculture* (33), 44-48
 - Tocher, D.R., Bell, J.G., Dick, J.R. Crampton, V.O., 2003. Effects of vegetable oil diets on Atlantic salmon hepatocyte desaturase activities and liver fatty acid compositions. *Lipids* (38), 723-732.
 - Tocher, D. R., Bendiksen, E.A., Campbell, P.J., Bell, J.G., 2008. The role of phospholipids in nutrition and metabolism of teleost fish. *Aquaculture* (280), 21-34.
 - Wold, P.A., Hoehne-Reitan, K., Cahu, C.L., Zambonino Infante, J., Rainuzzo, J., Kj-rsvik, E., 2007. Phospholipids vs. neutral lipids: effects on digestive enzymes in Atlantic cod (*Gadus morhua*) larvae. *Aquaculture* (272), 502-513.

The Effect of Soybean and Egg Lecithin on the Growth, Survival and Resistance to Anoxia Stress in Rainbow trout (*Oncorhynchus mykiss*) Alevin

H. Mohammadi Azarm¹, A. Abedian Kenari*¹ and M. Hedayati²

¹ Department of Fisheries, Faculty of Natural Resources and Marine Science, Tarbiat Modares University, I.R. Iran

² Institute of Endocrinology and Metabolism, Shaheed Beheshti University of Medical Sciences, I.R. Iran

(Received: 26/07/2011 , Accepted: 09/01/2012)

Abstract

This study was conducted to evaluate the effect of different lecithin sources and their levels on growth performance and resistance to anoxia stress of rainbow trout (*Oncorhynchus mykiss*) alevin with initial weight of 120 ± 4.08 g for 40 days. Soybean and egg lecithin at 2, 4 and 6 % levels were added to the diets. Seven treatments by three replications were used in this experiment. Therefore, 165 alevin were used per each 90 L tank. Alevin were fed to visual satiation at six meals per day for 40 days. At the end of experiment, alevins were fed by 2% egg lecithin had significantly higher ($p < 0.05$) growth performance compared to control group. Growth parameters in soybean lecithin groups were not significantly different compared to control group. Alevins were fed by 6 % egg lecithin had significantly higher survival rate than other treatments and control group. Also alevins were fed by all levels of egg lecithin had significantly higher resistance to anoxia stress compared to control group. The result of this experiment showed that egg lecithin induced to significantly affect on the growth performance and resistance to anoxia stress. The optimal dietary level of egg lecithin for this size is 2 % of diet.

Keywords: Soybean lecithin, Egg lecithin, Alevin, Growth, Stress, Rainbow trout