```
(MLC)
)
                                     // :
                                                     // :
```

Minnaert constant - "

Maximum Likelihood Classification - ٤

Topographic Normalization -°

108 .()) .() .() ' - Orientation of the Sun (solar azimuth)
Y - Solar elevation angle
F - Nadir view • - Off-nadir view

```
K
                         Cos i
\cos i = \cos(90 \quad \theta_s) \cos \theta_n + \sin(90 - \theta_s)
\sin \theta_n \cos(\phi_s - \phi_n)
                                                        ()
                                                                                 ) (Path-Row)
                                              \theta_{s}
                                                                                                           1
                                                                                                                   RMS
                                                                                                              (RGB)
                                      \mathbf{K}
                                             ()
\log(BV_{observed\lambda}\cos\theta) = \log BV_{normal\lambda} +
                                                                                                                         (RGB)
K log(cos i cos e)
                                                   ()
                                                                                                               :(
                                                                                                                        )
                                                                BV_{normal\lambda} = (BV_{observeed\lambda} \cos e) / (\cos^k i \cos^k e)
                                                                                                       BV_{normal\lambda}
                                      (GIS)
                                                                       Cos e
                                        (DEM)
```

⁻ Surface Roughness

⁴ - Digital Elevation Model

^{` -} Nearest Neighbor

^{*} Root Mean Squar

110 (cos i) (MLC) .()

¹ - Boolean logic

^{&#}x27; - Separability
' - Error matrix
' - Kappa Index of Agreement

.

.() ()

. ()

·

.

112

. .

TM1	TM2	TM3	TM4	TM5	TM7
1	1	1	1	1	1

TM1					
TM2					
11712					
TM3					
11013					
TMA					
TM4					
TMC					
TM5					
TM7					

...

·

'-Noise

... 116

.

	1	•					
TM1							
TM2							
TM3							
TM4							
TM5							
TM7							

TM1				
TM2				
TM3				
	1			
TM4				
TD 45				
TM5				
T) 47				
TM7				

... 118

				•		
					/	/
					,	,
					1	/
			/		1	
			,		,	1
			/			/
Г	I				,	,
					1	1
					1	1
					1	1
					1	1
					1	1
		1	1	1	1	1
					1	1

.

.

.

(Cosi)

) /) .(/ () (DEM)

' - Line-of-sight

() ()

- 1-Colby , J. D., 1991. Topographic Normalization in Rugged Terrain, Photogrammetric Engineering & Remote Sensing, 57(5): 531-537.
- 2-Congalton, R., 1991. A review of assessing the accuracy of classifications of remotely sensed data, Remote Sensing of Environment, Vol. 37:35-46.
- 3-ERDAS IMAGINE, 1996. ERDAS field guides, 3rd edition. ERDAS, Inc. Atlanta, Georgia. 4-Hodgson, M. E., & B.M. Shelley, 1993. Removing the topographic effect in remotely sensed. Imagery. ERDAS Monitor, Fall 1993. Contact Dr. Hodgson, Dept of Geography, University of Colorado, Boulder, CO 80309-0260.
- 5-Holben, B. N., & C.O. Justice, 1980. The Topographic effect on spectra response from Nadir points, Photogrammetric Engineering & Remote Sensing, Vol. 46(9): 1191-1200.
- 6-Jones, A. R., J. J. Settle, & B.K. Wyatt, 1988. Use of digital terrain data in the interpretation of SPOT-IHRV Multispectral imagery. International Journal of Remote Sensing. Vol. 9 (4): 669-682.
- 7-Kawata, Y.S. U., & T.Kusaka,1988. Radiometric correction for atmospheric and topographic effects on Landsat MSS images. International Journal of Remote Sensing, Vol. 9(4): 729-748.
- 8-Leprieur, C.E., J.M. Durand, & J.L. Peyron,1988. Influence of topogrphy on forest reflectance using Landsat Thematic Mapper and Digital Terrain data. Photogrammetric Engineering & Remote Sensing, Vol. 54(4):491-496.
- 9-MERA, 1996. MARS and Environment Related Applications. PHARE program, Forest Ecosystem Mapping., Final report, FOMI Remote Sensing Center, Budapest Hungray.
- 10-Pinter, P.J., R.D. Jackson, S.B. Idso, & R.J. Reginato, R.J. 1983. Diurnal patterns of wheat spectral reflectances. IEEE Transactions on Geoscience and Remote Sensing, GE-21, 156-163.

¹¹⁻ Proy, C., D. Tanre, & P.Y. Deschamps, 1989. Evaluation of topographic effects in remotely sensed data. Remote Sensing of Environment, Vol. 30:21-32. 12-Smith, J. A., T. L. Lin, & K.J. Ranson, 1980. The Lambertian assumption and Landsat Data. Photogrammetric Engineering & Remote Sensing Vol. 46:1183-89.

Studying the Effects of Topographic Normalization on Satellite Imageries Using Minnaert Constant

A.Sepehry¹

Abstract

Vegetation mapping is one of the important aims phytosociologists are after. Everincreasing improvement of satellite images in terms of their spatial and spectral resolution made them suitable for vegetation mapping. Satellites capture electromagnetic radiation reflected or emitted by objects on the earth. Electromagnetic radiation itself is a function of spectral property of the object, sun position (sun azimuth and elevation), and surface terrain. As a result, a given plant cover on different aspects of the terrain may be detected differently. This is especially true for images from mountainous environments. Thus, it is necessary to use techniques that can improve the amount of information obtained from such images. Several such techniques have evolved over the years, including the odeling of illumination effects using a DEM² and the calculation of surface roughness index such as Minnaert constant to adjust over correction caused by non lambertian models. This research attempts to apply such method to study the effects of topographic normalization on classification accuracy of Landsat TM image acquired for the preserved mountainous area of Jahan-nama in province of Golestan. The study area was classified using MLC³ method on images, which had been topographically normalized by applying Minnaert constant. The results of comparing topographically normalized thematic map with original thematic map show that the topographic normalization method applied over corrects the image, and that the overall classification accuracy on corrected image is lower than the uncorrected one. The over-correction seems to be due to the quality of DEM used in the procedure. Using the information about the sun position which is available from the Landsat header file, is probably the other source of over-correction. This is because it is the position of sun relative to the center of the image while the study area was located at the extreme end of the image. The research suggests that calculating the position of sun for all pixels involved may help deriving better results.

Key words: Topographic normalization, Minnaert constant, Landsat, TM, Vegetation, Classification, Jahan nama, Golestan

 $^{^{1}}$ - Asst. Prof., Faculty of Watershed and Range Management, Agricultural Seciences and Natural Resources University of Gorgan

² Digital Elevation Model

³ Maximum Likelihood Classification